



Characterization of CO Thermochemistry in Incident Shockwaves

Brett A. Cruden, Aaron M. Brandis
AMA, Inc

Megan E. MacDonald
Jacobs Engineering

AIAA AVIATION 2018
Atlanta, GA




- Motivation
- Features of CO Shocks
- Temperature vs Position from shock
 - Determination of CO Dissociation Rate
- C₂ Swan Band Radiation
 - Influence of C₂ reaction rates
- CO VUV and IR Radiation
- Summary

Motivation




- Spacecraft to Mars typically enter the atmosphere at 5-7 km/s
- Mars Atmosphere is 96% CO₂, 1.9% N₂, 1.9% Ar
- Radiative and Convective Heating Depend Upon Reactions in Shock Layer
- Most rates in use based upon old (60s-70s) shock tube studies
- Recent updates (i.e. Johnston 2014) based on data in mixture



Simplified Chemistries



- Extensive validation work has been performed on full chemistry set
- Model revisions* have also been based on measurements of full chemistry
- Studies of simplified chemistry sets may result in better informed modeling choices




* Johnston, C. O., and Brandis, A. M., "Modeling of nonequilibrium CO Fourth-Positive and CN Violet emission in CO₂-N₂ gases," *Journal of Quantitative Spectroscopy and Radiative Transfer*, Vol. 149, 2014, pp. 303-317

Simplified Chemistries





- Substituting CO for CO₂ minimizes the influence of CO₂ reactivity on results

• ~~Mars~~
CO





Start with CO

Electric Arc Shock Tube (EAST)




- Experiments Conducted in EAST Facility to study shock-heated CO
- 4" Aluminum Tube, 7.9 m from primary diaphragm to test section
- Initial condition of 0.1 and 0.25 Torr CO, shock velocities from 3-9 km/s
- Diagnostics
 - Tunable Diode Laser Spectroscopy (Mid-Wave Infrared)
 - Imaging Emission Spectroscopy (4 spectrometers covering VUV through mid-Infrared)

Analysis of CO Data





- TDLAS data*:
 - Measures one molecular line vs. time
 - Obtain translational temperature, CO number density
- Emission spectroscopy
 - Measure radiance versus position and wavelength over broad spectral range:
 - VUV (145-195nm, CO A-X transition), UV (190-330 nm), Visible (480-890 nm, C₂ Swan Bands), mid-Infrared (4000-5500 nm, CO vibrational)
 - Radiant Power depends on temperature, (excited) species number densities

* Macdonald, et al., 2:30 pm Thur.




Characteristics of CO Shocks

Equilibrium CO Shocks





- Temperatures from 4000-8000K
- Major species CO, C and O
- C₂ at 10-200 ppm, starting at 3.5 km/s
- C⁺ ion at 100 ppm at 4 km/s, 0.1% at 6 km/s, 1% at 10 km/s
- CO₂, CO⁺, O⁺ and O₂ are more minor species
- Mole fractions, temperatures not strong function of pressure from 0.1-0.25 Torr



- CO Dissociation



- CO Exchange



- O₂/C₂ Dissociation



- Ionization kicks in at higher velocity
- 5 sp, 5 rxn

Space of Valid Solutions



- 1D Shock must satisfy conservation equations:

Atom conservation:

$$n_{CO} + n_C + 2n_{C_2} = \frac{\rho_0 v_0}{M_{CO} v}$$

Stoichiometry:

$$n_C + 2n_{C_2} = n_O + 2n_{O_2}$$

(Atom + Stoichiometry conservation enforces Conservation of Mass)



Momentum:


$$\sum n_i (RT + M_i v^2) = p_0 + \rho_0 v_0^2$$

Energy:

$$\sum n_i \left(h_i(T) + M_i \left(\frac{1}{2} v^2 - h_0 - \frac{1}{2} v_0^2 \right) \right) = 0$$

- 4 equations, 7 unknowns (v , T , 5 n_i 's) – 3 DOF







Reaction Rate Analysis

CO Reaction Rates




- Reaction Rate may be inferred from T vs. x:





- $\omega = f(T)$ ← Mass fraction is a (narrowly bounded) function of Temperature, per conservation equations
- $\frac{d\omega_{CO}}{dx} = \frac{M_{CO}r_{CO}}{\rho_0 v_0}$ ← Derivative of mass fraction proportional to reaction rate
- $\frac{r_{CO}}{n_{CO}n_M} \approx k_{diss}$ ← Reaction rate dominated by CO dissociation by heavy particle collision

CO Dissociation Rate




- Reaction rate measured over many tests cluster around the rate curve reported by Hanson
 - Note that reaction at high Temperature is faster than the time scale of TDLAS : fit is less reliable/more scattered
 - Rate has large Arrhenius ($T^{-5.5}$) coefficient : suggests compound mechanism

Compound Mechanism for CO dissociation



$$k_{eff} = \frac{k_{diss}k_{exc}}{k_{de-exc} + k_{diss}}$$

$$= \frac{k_{diss}k_{exc}}{\left(\frac{n_{CO^*}}{n_{CO(X)}}\right)_{eq} k_{exc} + k_{diss}}$$





- Compound mechanism involves excitation to an excited state that then dissociates
 - At high temperature, excitation is rate limiting step, excited state is populated *below* equilibrium
 - At low temperature, dissociation is rate limiting, excited state is equilibrated
- What is the excited state?
 - Some literature has suggested M not inert and could be intermediate such as CO₂, C₂, O₂
 - An obvious choice may be the CO metastable, CO(a)


Intermediate States

- Evaluation of rate using a, a', A, d, e states of CO all can plausibly explain data

$$k_{exc} = A_{exc} T^{1/2} \exp\left(-\frac{E^*}{T}\right) \quad k_{diss} = A_{diss} T^{1/2} \exp\left(-\frac{E - E^*}{T}\right)$$






- Dissociation rates derived from a, a' are within an order of magnitude of estimates from Park
- De-excitation rates are somewhat lower than rates measured at 300K
- Likely a combination of the above



Data Comparisons




Temperature

Temperature Predictions 3-6 km/s




- At 3.4 km/s – No dissociation, trend driven by T-Tv relaxation
- At 4.4 km/s and 5.7 km/s
 - Schwenke and Johnston Rate too slow
 - Hanson rate matches data
 - T_v trend looks ok

Temperature Predictions 6-9 km/s



- Schwenke Rate too slow
- At 6.6-7.5 km/s
 - Johnston and Hanson rates similarly match data
- At 8.6 km/s
 - Data relaxes faster than any rate predicts





C_2 Thermochemistry

C₂ Rates



- There are two rates that matter for C₂ radiation : Dissociation and Exchange





- There is about 1 OOM difference between Park and literature for dissociation rate
- Up to 2 OOM difference between Park and literature for exchange rate

C₂ Rate Sensitivity




5.6 km/s






- Radiation shows significant sensitivity to exchange
 - Park's exchange puts peak at wrong location, makes it wider
- Result is less sensitive to dissociation rate
 - Discrepancy at the peak may be due to Boltzmann model
 - If not, rate would need to be even faster
- Fairbarn's rates appear more consistent with data

C₂ Radiance Predictions






- Except for peak, agreement is very good
- C₂ reaction rates from Fairbarn agree better than those of Park
- Further improvement would be construction of a non-Boltzmann model

C₂ Radiance – 4-7 km/s




- 4.4 and 5.7 km/s
 - Schwenke overpredicts (T too high)
 - Hanson+Fairbarn overpredicts peak, good elsewhere
 - Johnston (Park) gets trend wrong
- 6.6 km/s
 - Similar observations, Johnston showing better agreement

C₂ Radiance – 7-9 km/s






- Johnston agrees well with data at these velocities
 - Boltzmann model for C₂
- Hanson/Fairbarn relaxation trend is no longer matched at 8.6 and 9.5 km/s



CO Radiation Analysis




VUV and mid-IR

CO VUV Radiance – 4-7 km/s






- 4.4 and 5.7 km/s
 - Schwenke overpredicts (T too high)
 - Hanson+Fairbarn matches data
 - Johnston over/under
- 6.6 km/s
 - Hanson and Johnston same result

CO VUV Radiance – 7-9 km/s




- Johnston agrees well with data at these velocities
- Hanson overprediction becomes worse as velocity increases

VUV Spectral Comparison (5.7 km/s)





- VUV Prediction does not simultaneously match low and high wavelength region




Fitting of CO VUV Bands



- Fits of the VUV spectrum:
 - Requires 3 Temperatures in Non-equilibrium
 - Result in elevated CO density (compared to what conservation allows)






CO MWIR Radiance – 3-6 km/s






- CO IR radiance not very sensitive to reaction rates in this velocity regime
- Always underpredicted

CO MWIR Radiance – 6-9 km/s






- Slower reaction rates predict overshoots that are not observed in experiment
- Still underpredicted

IR Underprediction





- Underprediction nearly constant with wavelength
- Correction required would be $\sim 1.3x$

Absorbance is Overpredicted



- Absorbance overpredicted by ~25%
 - Opposite to trend in Emission, cannot be error in linestrength or CO number density
- At 8.6 km/s, temperature prediction does not match


CO MWIR Predictions



- Different linelists have been tested in NEQAIR and all produce substantially the same result
- Alberti, et al. (2017) showed that HITRAN underpredicts experimental data above ~ 1200 K at pressures up to 3 bar
 - Consistent with emission trend
- To underpredict the band but overpredict a line indicates there are additional contributors to the band
 - Additional excited states of CO not accounted for?
 - May have implications for CO partition function and thermodynamics

- Analysis of Temperature Relaxation Finds CO dissociation rate in agreement with that of Hanson
 - It is suggested this is a compound reaction that proceeds through CO metastable levels
 - Excited states must be considered in QCT calculations!
 - CO Metastable is at about half of the dissociation energy
- Emission trends are consistent with CO rate of Hanson below ~ 7 km/s, and Johnston above 7 km/s
- C₂ Emission more consistent with rates of Fairbarn than Park

- Deviation of Data at High Velocity
 - Role of electron impact dissociation of CO?
 - Need for merged Johnston/Hanson reaction rate?
- Overprediction of C₂ Swan band at shock front
 - Non-Boltzmann modeling of C₂
- Inconsistencies in CO radiance
 - Absorption and emission disagree with predictions in opposite direction
 - HITRAN database shown not to match high temperature CO data
 - Possible errors in CO Partition function and energy levels?
 - Predicted shape of CO 4th Positive radiation
 - Dipole moments and potential energy surfaces for CO




Backup


Rate Extracted – Assuming exchange



O atom exchange



C atom exchange



- Rates fail to collapse – not correlated with O or C atom fraction
- Rates inconsistent with predicted rates
- Suggests exchange does not drive dissociation